

IOWA STATE UNIVERSITY

Digital Repository

Integrated Crop Management News

Agriculture and Natural Resources

2-12-2007

Allelopathy: A cause for yield penalties in corn following corn?

Roger W. Elmore

Iowa State University, relmore@iastate.edu

Lori Abendroth

Iowa State University, labend@iastate.edu

Follow this and additional works at: <http://lib.dr.iastate.edu/cropnews>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), and the [Agronomy and Crop Sciences Commons](#)

Recommended Citation

Elmore, Roger W. and Abendroth, Lori, "Allelopathy: A cause for yield penalties in corn following corn?" (2007). *Integrated Crop Management News*. Paper 1158.

<http://lib.dr.iastate.edu/cropnews/1158>

This Article is brought to you for free and open access by the Agriculture and Natural Resources at Digital Repository @ Iowa State University. It has been accepted for inclusion in Integrated Crop Management News by an authorized administrator of Digital Repository @ Iowa State University. For more information, please contact digirep@iastate.edu.

INTEGRATED CROP MANAGEMENT

Get the latest research-based information on crops. [Sign up to be notified](#) when new content is available!

ICM > 2007 > IC-498 (1) -- February 12, 2007

Current Newsletter

You are viewing **archives** for the newsletter from 1993-2007. For current news, see [Integrated Crop Management News](#).

Archives 1993-2007



Announcements



Crop Production



Insects and Mites



Pesticide Education



Plant Diseases



Soils



Weed Management

[Image Gallery](#)

Printable Version

Printable version of this page

Related Articles

Conservation systems: Challenges and benefits
October 1, 2007

Nutrient removal when harvesting corn stover
August 6, 2007

How residue removal affects nutrient cycling
May 21, 2007

Residue removal and potential environmental

Allelopathy: A cause for yield penalties in corn following corn?

by Roger Elmore and Lori Abendroth, Department of Agronomy

Based on research and in-field experience, we are confident yields are reduced when corn follows corn in Iowa cropping systems. The question many ask though is why? What are the causes? Some point to allelopathy and wonder if this is the major cause.

Researchers have examined this specific question over the last two decades. We will discuss some of their research findings in this article.

According to Webster's dictionary, allelopathy is defined as the suppression of growth of one plant species by another plant species, due to the release of toxic substances called allelochemicals. Many crops are allelopathic when grown with other crops or when they are grown sequentially. Allelochemicals are released through many processes, including leaching by precipitation, decomposition of crop residues by microbial activity, and through root exudates. Allelopathy was documented in the 1830s; yet even at that time, scientists knew its impact was minimized with crop rotation.

Autotoxicity is a specific type of allelopathy.

Autotoxicity occurs when the allelochemicals released from a specific crop affect that same crop planted at a later time. Corn is one of several autotoxic crops. Other plants are only allelopathic to differing plant species--but not to themselves.

Research conducted in a laboratory at Iowa State University showed that corn seeds placed into corn residue taken out of the field limited seedling dry root and shoot weight. Yet, when the residue was decomposed (as occurs prior to spring) the phytotoxins were inactivated by microbial breakdown, thereby not affecting plant growth. In another laboratory study by Ohio State University, corn seeds incubated in corn residue extracts had a 60 percent reduction in germination compared to the control, whereas seed incubated in soybean residue extracts had a 40 percent reduction in germination. Coleoptile (shoot), radicle (primary root), and secondary root lengths were all reduced more by corn residue than soybean residue.

Do these laboratory results, which show some reduction in seed germination and growth, apply to actual field situations?

Corn seedlings emerged sooner in soybean residue than those in corn residue in Iowa State University field research.

Tillage systems (no-till and mold board plow), residue types (corn or soybean), and removal

consequences
April 23, 2007

Disease management in
corn-following-corn
fields

February 12, 2007

Tillage challenges in
managing continuous
corn

February 12, 2007

Equipment
considerations for
producing corn after corn

February 12, 2007

Can residue be
managed successfully
with no-till?

May 15, 2006

Why conservation
systems are the right
choice this fall

October 10, 2005

Residue management
and manure application
October 21, 2002

schemes (residue removed from seed row in bands of 0, 3.14, 6.3, 12.6, and 30 inches wide) were evaluated in this experiment. Removing residue from the top of the row had greater effects on corn growth and yield than either tillage or residue type. They concluded that several soil characteristics were not reducing yield, including allelopathy.




Residue is not the primary cause of yield penalties in corn following corn but may be a contributing factor among many. (Lori Abendroth)

The factor that was limiting yield and plant growth was the residue's effect on soil temperature. Removing a 6-inch band of residue over the planting row significantly limited the yield-reducing effect of no-till, and any inherent yield reductions due to allelopathy from the residue. Scientists who studied in-row residue placement in no-till systems in Guelph, Canada, did not conclude this however. They found that reducing the amount of in-row residue did not always overcome the "yield-limiting effects of no-till corn production."

Obviously, there is variability in these research reports, yet we can draw some conclusions concerning the toxicity of corn residue to the following corn crop. Corn residue appears to negatively impact germination and early seedling growth. This is verified by the laboratory research and the first field report. Laboratory research could single out the impact of residue without having temperature as an influencing factor; field research typically cannot do this. Therefore, in field research, we cannot separate the influence of residue on overall soil temperature and seedbed conditions. In the Iowa research, the movement of residue away from the row reduced the negative impact of the residue. Yet, in cooler northern soils (such as Guelph, Canada), moving the residue did not eliminate some of the negative impacts from the residue.

Allelopathy appears to be of some influence in corn seed germination and early season growth but is greatly minimized when the residue is moved away from the row. Therefore, it is not likely to be the primary cause of yield penalties in corn following corn but, instead, is a contributing factor among many.

Proportional acreage by crop—Iowa 1926–2006



Source: United States Department of Agriculture.

Find links to research articles discussed here at our [Web site](#) under "Cropping Systems."

Roger Elmore is professor of agronomy with research and extension responsibilities in corn production. Lori Abendroth is an agronomy specialist with research and extension responsibilities in corn production.

This article originally appeared on pages 16-17 of the IC-498 (1) -- February 12, 2007 issue.

Updated 02/13/2007 - 3:13pm